

ON THE FUTAKI INVARIANTS OF COMPLETE INTERSECTIONS

ZHIQIN LU

1. INTRODUCTIONS

In 1983, Futaki [2] introduced his invariants which generalize the obstruction of Kazdan-Warner to prescribe Gauss curvature on S^2 . The Futaki invariants are defined for any compact Kähler manifold with positive first Chern class that has nontrivial holomorphic vector fields. Their vanishing are necessary conditions to the existence of Kähler-Einstein metric on the underlying manifold.

Let M be a compact Kähler manifold with positive first Chern class $c_1(M) > 0$. Choosing an arbitrary positive $(1, 1)$ form ω in $c_1(M)$ as a Kähler metric on M , we can find a smooth function f on M , determined up to a constant, such that the following

$$(1.1) \quad Ric(\omega) - \omega = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} f$$

holds. Let $\mathfrak{b}(M)$ be the Lie algebra of holomorphic vector fields on M . The Futaki invariants are defined as

$$F : \mathfrak{b}(M) \rightarrow C, F(X) = \int_M X(f) \omega^n.$$

Ding and Tian [1] took a further step in introducing the Futaki invariants to Fano normal varieties. This is not only a generalization of Futaki invariants to singular varieties, but also has important application in Kähler-Einstein geometry. In [8], the generalized K-energy on normal varieties was introduced and a stability criteria for the hypersurface or complete intersection was established by using the notion of the generalized K-energy. The Futaki invariants on singular varieties are related to the stability of Fano manifolds due to the work of Tian [9]. To be more precise, checking the K -stability of a Fano manifold is the same as checking the sign of the real part of the Futaki

Date: June 3rd, 1998.

1991 Mathematics Subject Classification. Primary: 32F07; Secondary: 58G03.

Key words and phrases. Futaki invariant, complete intersection, Kähler-Einstein metric.

invariants on the degenerations of the Fano manifold. Because of this, we need an effective way to compute the Futaki invariants on singular varieties.

In this paper, we give a simple formula for the Futaki invariants of Fano complete intersections. The main theorem of this paper is:

Theorem 1.1. *Let M be the $N - s$ dimensional normal Fano variety in CP^N defined by the homogeneous polynomials $F_1, \dots, F_s$ of degree $d_1, \dots, d_s$ respectively. Let X be a holomorphic vector field on CP^N such that*

$$(1.2) \quad XF_i = \kappa_i F_i, \quad i = 1, \dots, s$$

for constants $\kappa_1, \dots, \kappa_s$. Then the Futaki invariant $F(X)$ is

$$(1.3) \quad F(X) = m^{N-s} \prod_{i=1}^s d_i \left(- \sum_{i=1}^s \kappa_i + \frac{m}{N-s+1} \sum_{i=1}^s \frac{\kappa_i}{d_i} \right),$$

where $m = N + 1 - d_1 - \dots - d_s$.

Corollary 1.1. *If M is a hypersurface in CP^N defined by the homogeneous polynomial F of degree d and if $XF = \kappa F$, then*

$$F(X) = -(N+1-d)^{N-1} \frac{(N+1)(d-1)}{N} \kappa.$$

In particular, $Re F(X)$ and $-Re \kappa$ have the same sign.

The formula is new even in the case when M is a hypersurface or an orbifold. If the zero locus of the holomorphic vector field X on M is a smooth manifold, then using the residue formula of the Atiyah-Bott-Lefschetz type, Futaki was able to develop a method to compute his invariants by the information of the vector field X and the manifold M near the zero locus of the vector field [3]. In [1], the authors developed the method to compute the Futaki invariants on 2-dimensional Kähler orbifolds. In [6], the Futaki invariants for toric varieties were calculated.

Acknowledgment. The author thanks G. Tian for his mathematical insights and encouragements during the preparation of this paper. He also thanks D. Phong, M. Kuranishi, H. Pinkham, L. Borisov and Z. Wu for the discussions of this topic.

2. PRELIMINARIES

Let $X = (a_{ij})_{\substack{1 \leq i \leq N \\ 1 \leq j \leq N}} \in sl(N+1, C)$ be a traceless $(N+1) \times (N+1)$ matrix. X defines a holomorphic vector field

$$(2.1) \quad X = \sum_{i,j=0}^N a_{ij} z^i \frac{\partial}{\partial z^j}$$

on C^{N+1} and a smooth function

$$(2.2) \quad \theta = -X \log \sum_{i=0}^N |z^i|^2$$

on $C^{N+1} - \{0\}$, where $(z^0, \dots, z^N)$ is the coordinates of C^{N+1} .

Both X and θ descends to a vector field and a smooth function on the projective space CP^N , respectively. Let ω_{FS} be the Fubini-Study metric of CP^N . Then we have the following relation:

$$(2.3) \quad i(X)\omega_{FS} = -\frac{\sqrt{-1}}{2\pi} \bar{\partial}\theta.$$

Suppose M is an n -dimensional Q -Fano normal variety in CP^N and suppose that $\omega = \alpha\omega_{FS} \in c_1(M)$ for a constant $\alpha > 0$. If X is a vector field on CP^N such that the one parameter group generated by the real part of X leaves M invariant, we say that X is tangent to M . Suppose $(x^1, \dots, x^n)$ is the local holomorphic coordinates at some smooth point p of M . Equation (2.3) can be written as

$$(2.4) \quad X^i = -\alpha g^{i\bar{j}} \frac{\partial \theta}{\partial \bar{x}^i}, \quad i = 1, \dots, n, \quad X = \sum_{i=1}^n X^i \frac{\partial}{\partial x^i},$$

where $(g^{i\bar{j}})$ is the inverse matrix of $(g_{i\bar{j}})$ and $(g_{i\bar{j}})$ is the metric matrix of ω .

We define the divergence of X on M by

$$(2.5) \quad \text{div } X = \frac{\partial X^i}{\partial x^i} + \sum_{i=1}^n X^i \frac{\partial}{\partial x^i} \log \det(g_{i\bar{j}}).$$

The following lemma is the observation on which the whole paper is based.

Lemma 2.1 ([9]). *If M is a normal projective variety, then*

$$(2.6) \quad \text{div } X - \alpha\theta + X(f) = \text{const}$$

where the function f is defined as

$$(2.7) \quad Ric(\omega) - \omega = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} f$$

Proof: A straightforward computation yields

$$\frac{\sqrt{-1}}{2\pi} \bar{\partial} \operatorname{div} X = -i(X) Ric(\omega).$$

By Equation (2.7), we see that

$$\frac{\sqrt{-1}}{2\pi} \bar{\partial} \operatorname{div} X = -i(X)\omega - \frac{\sqrt{-1}}{2\pi} i(X) \partial \bar{\partial} f = \frac{\sqrt{-1}}{2\pi} \alpha \bar{\partial} \theta - \frac{\sqrt{-1}}{2\pi} \bar{\partial} X(f).$$

Thus $\operatorname{div} X - \alpha \theta + X(f)$ is a holomorphic function on the normal variety M which must be a constant. □

Corollary 2.1. *The Futaki invariant can be written as*

$$F(X) = \operatorname{const} \operatorname{vol}(M) + \alpha \int_M \theta \omega^n.$$
□

3. AN EXPLICIT EXPRESSION OF THE FUNCTION f

Suppose M is a complete intersection of CP^N . That is, M is the zero locus of homogeneous polynomials $F_1, \dots, F_s$ in CP^N with degree $d_1, \dots, d_s$, respectively and the dimension of M is $N - s$. By the adjunction formula, the anticanonical bundle of M is

$$K_M^{-1} = (N + 1 - d_1 - \dots - d_s)H,$$

where H is the hyperplane bundle of CP^N . We assume that M is a normal variety. There is a unique function f (up to a constant), defined on the regular part M_{reg} of M , such that if $\omega = (N + 1 - d_1 - \dots - d_s)\omega_{FS}|_M$, then

$$Ric(\omega) - \omega = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} f \quad \text{on } M_{reg}.$$

In this section, we write out the above function f explicitly. The idea is to trace the proof of the well known adjunction formula. But here we work on the metric level rather than the cohomological level. This makes the notations a little bit complicated.

We begin by the following general setting: Let V be a Kähler manifold of dimension N and V_1 be a submanifold of dimension $N - s$ defined by holomorphic functions $G_1 = G_2 = \dots = G_s = 0$. Suppose U_1 is an open set of V such that

- (1) $(x^1, \dots, x^N)$ is a local holomorphic coordinate system of V ;
 (2) On U_1 , we have

$$\text{rank} \frac{\partial(G_1, \dots, G_s)}{\partial(x^1, \dots, x^s)} = s;$$

- (3) There are holomorphic functions $f_1, \dots, f_s$ on $U_1 \cap V_1$ such that

$$\begin{cases} x^1 = f_1(x^{s+1}, \dots, x^N) \\ x^2 = f_2(x^{s+1}, \dots, x^N) \\ \dots\dots\dots \\ x^s = f_s(x^{s+1}, \dots, x^N) \end{cases}.$$

In particular, $(x^{s+1}, \dots, x^N)$ is the local holomorphic coordinate system of $U_1 \cap V_1$.

Suppose $g_1 = \sum_{\substack{s+1 \leq i \leq N \\ s+1 \leq j \leq N}} g_{1i\bar{j}} dx^i \wedge d\bar{x}^j$ is the restriction of the Kähler metric of V on $U_1 \cap V_1$. Define $\det g_1 = \det(g_{1i\bar{j}})_{\substack{s+1 \leq i \leq N \\ s+1 \leq j \leq N}}$. Of course,

$\det g_1$ is not a global function on V_1 . In order to study the change of the $\det g_1$ with respect to the change of the local holomorphic coordinates, we assume that there is another neighborhood $(U_2, (y^1, \dots, y^N))$ of V such that $U_1 \cap U_2 \cap V_1 \neq \emptyset$. As before, we assume that

$$\text{rank} \frac{\partial(G_1, \dots, G_s)}{\partial(y^1, \dots, y^s)} = s$$

and on $U_2 \cap V_1$, we have

$$\begin{cases} y^1 = g_1(y^{s+1}, \dots, y^N) \\ y^2 = g_2(y^{s+1}, \dots, y^N) \\ \dots\dots\dots \\ y^s = g_s(y^{s+1}, \dots, y^N) \end{cases}$$

for holomorphic functions $g_1, \dots, g_s$ on $U_2 \cap V_1$. $(y^{s+1}, \dots, y^N)$ is the local holomorphic coordinate system of $U_2 \cap V_1$. Let

$$g_2 = \sum_{\substack{s+1 \leq i \leq N \\ s+1 \leq j \leq N}} g_{2i\bar{j}} dy^i \wedge d\bar{y}^j$$

be the restriction of the Kähler metric of V on $U_2 \cap V_1$. Define $\det g_2 = \det(g_{2i\bar{j}})_{\substack{s+1 \leq i \leq N \\ s+1 \leq j \leq N}}$. Then we have

Proposition 3.1 (Adjunction Formula). *With the above notations, on $V_1 \cap U_1 \cap U_2 \neq \emptyset$, we have*

$$\det(g_1) = \det(g_2) \left| \det \left(\frac{\partial y^j}{\partial x^i} \right)_{\substack{1 \leq i \leq N \\ 1 \leq j \leq N}} \right|^2 \cdot \frac{\left| \det \frac{\partial(G_1, \dots, G_s)}{\partial(y^1, \dots, y^s)} \right|^2}{\left| \det \frac{\partial(G_1, \dots, G_s)}{\partial(x^1, \dots, x^s)} \right|^2}.$$

Proof: Let

$$\begin{cases} \tilde{x}^i = x^i - f_i(x^{s+1}, \dots, x^N) & 1 \leq i \leq s \\ \tilde{x}^i = x^i & i > s \end{cases},$$

and

$$\begin{cases} \tilde{y}^j = y^j - g_j(y^{s+1}, \dots, y^N) & 1 \leq j \leq s \\ \tilde{y}^j = y^j & j > s \end{cases}.$$

Then V_1 is locally defined by $\tilde{x}^i = 0$ ($1 \leq i \leq s$) or $\tilde{y}^j = 0$ ($1 \leq j \leq s$). In particular, on V_1 we have

$$(3.1) \quad \frac{\partial \tilde{x}^i}{\partial \tilde{y}^j} = 0, \quad 1 \leq i \leq s, \quad s+1 \leq j \leq N.$$

Before going further, we make the following conventions:

- (1) $\det\left(\frac{\partial x^i}{\partial y^j}\right)_{\substack{1 \leq i \leq N \\ 1 \leq j \leq N}}$ and $\det\left(\frac{\partial \tilde{x}^i}{\partial \tilde{y}^j}\right)_{\substack{1 \leq i \leq N \\ 1 \leq j \leq N}}$ denote the restriction to V_1

of the Jacobi determinant of the transform $(y^1, \dots, y^N) \rightarrow (x^1, \dots, x^N)$ and $(\tilde{y}^1, \dots, \tilde{y}^N) \rightarrow (\tilde{x}^1, \dots, \tilde{x}^N)$ respectively;

- (2) $\det\left(\frac{\partial \tilde{x}^i}{\partial \tilde{y}^j}\right)_{\substack{1 \leq i \leq s \\ 1 \leq j \leq s}}$ and $\det\left(\frac{\partial \tilde{x}^i}{\partial \tilde{y}^j}\right)_{\substack{s+1 \leq i \leq N \\ s+1 \leq j \leq N}}$ are the determinant of the

submatrices of $\left(\frac{\partial \tilde{x}^i}{\partial \tilde{y}^j}\right)_{\substack{1 \leq i \leq N \\ 1 \leq j \leq N}}$;

- (3) Since $(x^{s+1}, \dots, x^N)$ and $(y^{s+1}, \dots, y^N)$ are local coordinates of $U_1 \cap V_1$ and $U_2 \cap V_2$ respectively, there is the transform $(y^{s+1}, \dots, y^N) \rightarrow (x^{s+1}, \dots, x^N)$ by

$$x^i = x^i(g_1(y^{s+1}, \dots, y^N), \dots, g_s(y^{s+1}, \dots, y^N), y^{s+1}, \dots, y^N)$$

for $s+1 \leq i \leq N$ and $\det\left(\frac{\partial x^i}{\partial y^j}\right)_{\substack{s+1 \leq i \leq N \\ s+1 \leq j \leq N}}$ denotes the Jacobi

determinant of the above transform.

If $s+1 \leq i, j \leq N$, then

$$\frac{\partial \tilde{x}^i}{\partial \tilde{y}^j} = \frac{\partial x^i(g_1, \dots, g_s, y^{s+1}, \dots, y^N)}{\partial y^j}.$$

Using Equation (3.1), we have

$$(3.2) \quad \det\left(\frac{\partial x^i}{\partial y^j}\right)_{\substack{1 \leq i \leq N \\ 1 \leq j \leq N}} = \det\left(\frac{\partial \tilde{x}^i}{\partial \tilde{y}^j}\right)_{\substack{1 \leq i \leq s \\ 1 \leq j \leq s}} \cdot \det\left(\frac{\partial x^i}{\partial y^j}\right)_{\substack{s+1 \leq i \leq N \\ s+1 \leq j \leq N}}.$$

If $1 \leq i, j \leq s$, then

$$\frac{\partial \tilde{x}^i}{\partial \tilde{y}^j} = \sum_{k=1}^s \frac{\partial \tilde{x}^i}{\partial y^k} \cdot \frac{\partial y^k}{\partial \tilde{y}^j} = \frac{\partial x^i}{\partial y^j} - \sum_{k=s+1}^N \frac{\partial f_i}{\partial x^k} \cdot \frac{\partial x^k(g_1, \dots, g_s, y^{s+1}, \dots, y^N)}{\partial y^j}.$$

Thus we have

$$(3.3) \quad \sum_{i=1}^s \frac{\partial G_k}{\partial x^i} \cdot \frac{\partial \tilde{x}^i}{\partial \tilde{y}^j} = \frac{\partial G_k}{\partial y^j}.$$

The proposition follows from Equation (3.2) and (3.3). $\square$

We are going to use the above proposition in the case of complete intersections of CP^N . Since M is defined by the zero locus of homogeneous functions, we must make some necessary adjustment because homogeneous polynomials are *not* functions on CP^N .

Let U_α ($\alpha = 0, \dots, n$) be the standard covering of CP^N defined by $U_\alpha = \{Z_\alpha \neq 0\}$ where $[Z_0, \dots, Z_N]$ is the homogeneous coordinates of CP^N . Suppose $z_i^\alpha = Z_i/Z_\alpha$ ($i \neq \alpha$) be the standard coordinates on U_α . Let

$$F_i^\alpha(z_0^\alpha, \dots, z_\alpha^\alpha, \dots, z_N^\alpha) = F(Z_0, \dots, Z_N)/(Z_\alpha)^{d_i}, \quad 1 \leq i \leq s.$$

For each $\{\alpha, \beta_1, \dots, \beta_s\} \subset \{0, \dots, N\}$, define

$$U_{\alpha\beta_1\dots\beta_s} = \{p \in U_\alpha \mid \text{rank} \frac{\partial(F_1^\alpha, \dots, F_s^\alpha)}{\partial(z_{\beta_1}^\alpha, \dots, z_{\beta_s}^\alpha)} = s\}.$$

Then it is clear that $\cup U_{\alpha\beta_1\dots\beta_s} \supset M_{reg}$.

At each point $p \in M_{reg} \cap U_{\alpha\beta_1\dots\beta_s}$, $(z_\beta^\alpha, \beta \notin \{\alpha, \beta_1, \dots, \beta_s\})$ can be used as local coordinate system at p . Let $g_{\alpha\beta_1\dots\beta_s}$ be the corresponding metric matrix and let $\det g_{\alpha\beta_1\dots\beta_s}$ be its determinant. Define

$$(3.4) \quad \xi_{\alpha\beta_1\dots\beta_s} = \det(g_{\alpha\beta_1\dots\beta_s}) \left| \det \frac{\partial(F_1^\alpha, \dots, F_s^\alpha)}{\partial(z_{\beta_1}^\alpha, \dots, z_{\beta_s}^\alpha)} \right|^2 \left(1 + \sum_{i \neq \alpha} |z_i^\alpha|^2\right)^{N+1-d_1-\dots-d_s}.$$

Then we have

Lemma 3.1. $\{\xi_{\alpha\beta_1\dots\beta_s}\}$ defines a global positive function of M_{reg} .

Proof: A straightforward computation shows (cf. [4, pp. 146])

$$dz_0^{\alpha_2} \wedge \cdots \widehat{dz_{\alpha_2}^{\alpha_2}} \cdots \wedge dz_N^{\alpha_2} = \left(\frac{Z_{\alpha_1}}{Z_{\alpha_2}} \right)^{N+1} dz_0^{\alpha_1} \wedge \cdots \widehat{dz_{\alpha_1}^{\alpha_1}} \cdots \wedge dz_N^{\alpha_1}.$$

The lemma follows from Proposition 3.1, Equation (3.4) and the above equation. $\square$

Theorem 3.1. *Let f be the function on M_{reg} , defined by*

$$(3.5) \quad f = -\log \xi_{\alpha\beta_1 \cdots \beta_s} \quad \text{on} \quad U_{\alpha\beta_1 \cdots \beta_s}.$$

Then

$$\frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} f = Ric(\omega) - \omega,$$

where $\omega = (N + 1 - d_1 - \cdots - d_s)\omega_{FS}|_M$. $\square$

4. THE TRACE OF THE ACTION ON $F_1, \dots, F_s$

Let M be the variety defined in the previous section. The vector field $X = \sum_{i,j=0}^N a_{ij} Z_j \frac{\partial}{\partial Z_i}$ naturally acts on F_i by

$$(4.1) \quad XF_i = \sum_{i,j=0}^N a_{ij} Z_j \frac{\partial F_i}{\partial Z_i}, \quad i = 1, \dots, s.$$

Suppose $\mathcal{V}$ is the vector space spanned by $F_1, \dots, F_s$. Since X is tangent to M , X is an automorphism on $\mathcal{V}$.

The main result of this section is,

Theorem 4.1. *Let κ be the trace of the automorphism of X on $\mathcal{V}$. Then*

$$\operatorname{div} X + X(f) - (N + 1 - d_1 \cdots - d_s)\theta = -\kappa.$$

Proof: We adopt all notations from last section. Consider a smooth point p of M_{reg} in $U_{\alpha\beta_1 \cdots \beta_s}$. From Equation (2.2), the function θ in the local coordinates $\{z_j^\alpha, j \neq \alpha\}$ is

$$(4.2) \quad \theta = -\tilde{X} \log(1 + \sum_{i \neq \alpha} |z_i^\alpha|^2) - a_{\alpha\alpha} - \sum_{j \neq \alpha} a_{\alpha j} z_j^\alpha.$$

By (2.1), let

$$(4.3) \quad \tilde{X} = \sum_{i \neq \alpha} \left(\left(\sum_{j \neq \alpha} a_{ij} z_j^\alpha - a_{\alpha j} z_j^\alpha z_i^\alpha \right) + (a_{i\alpha} - a_{\alpha\alpha} z_i^\alpha) \right) \frac{\partial}{\partial z_i^\alpha}.$$

Let the holomorphic vector field X on M be written as

$$(4.4) \quad X = \sum_{i \notin \{\alpha, \beta_1, \dots, \beta_s\}} X^i \frac{\partial}{\partial z_i^\alpha}.$$

If $i : M \rightarrow CP^N$ is the embedding, then $i_*X = \tilde{X}$.

By Equation (3.4), (3.5), (4.2), (4.4) and the definition of $\operatorname{div} X$ in Equation (2.5), we see that

$$(4.5) \quad \begin{aligned} & \operatorname{div} X + X(f) - (N + 1 - d_1 - \dots - d_s)\theta \\ &= \sum_{i \notin \{\alpha, \beta_1, \dots, \beta_s\}} \frac{\partial X^i}{\partial z_i^\alpha} - X \log \det \frac{\partial(F_1^\alpha, \dots, F_s^\alpha)}{\partial(z_{\beta_1}^\alpha, \dots, z_{\beta_s}^\alpha)} \\ & \quad + (N + 1 - d_1 - \dots - d_s)(a_{\alpha\alpha} + \sum_{j \neq \alpha} a_{\alpha j} z_j^\alpha). \end{aligned}$$

Before going on, we need a general elementary lemma. To begin, we use the general setting on page 4. In addition, we let X be a holomorphic vector field of V such that X is tangent to V_1 . In what follows, we temporarily distinguish the X on V and the X on V_1 . So let's denote the X on V to be $\tilde{X}$. In the local coordinates, $\tilde{X}$ is

$$\tilde{X} = \sum_{i=1}^N \tilde{X}^i \frac{\partial}{\partial x^i}.$$

Then X on V_1 can be written as

$$X = \sum_{i=s+1}^N X^i \frac{\partial}{\partial x^i},$$

where from the chain rule,

$$(4.6) \quad X^i = \tilde{X}^i(f_1, \dots, f_s, X^{s+1}, \dots, X^N), \quad s+1 \leq i \leq N.$$

If $i : V_1 \rightarrow V$ is the embedding, then $i_*X = \tilde{X}$.

We have the following elementary lemma:

Lemma 4.1. *Let*

$$A = \frac{\partial(G_1, \dots, G_s)}{\partial(x^1, \dots, x^s)}$$

and let $A_{ij} = \frac{\partial G_i}{\partial x^j}$ for $i, j = 1, \dots, s$ and (A^{ij}) be the inverse matrix of A . Then on V_1 , we have

$$\sum_{i=s+1}^N \frac{\partial X^i}{\partial x^i} - \tilde{X} \log \det A = \sum_{i=1}^N \frac{\partial \tilde{X}^i}{\partial x^i} - \sum_{i,j=1}^s A^{ji} \frac{\partial}{\partial x^j} \tilde{X} G_i.$$

Proof: By definition,

$$\tilde{X} \log \det A = \sum_{k=1}^N \sum_{i,j=1}^s A^{ji} \tilde{X}^k \frac{\partial^2 G_i}{\partial x^j \partial x^k}.$$

We can write the above equation as

$$(4.7) \quad \tilde{X} \log \det A = \sum_{i,j=1}^s (A^{ji} \frac{\partial}{\partial x^j} \tilde{X} G_i - \sum_{k=1}^N A^{ji} \frac{\partial \tilde{X}^k}{\partial x^j} \frac{\partial G_i}{\partial x^k}).$$

By the implicit differentiation, we see that on V_1 ,

$$(4.8) \quad \frac{\partial f_j}{\partial x^k} = - \sum_{i=1}^s A^{ji} \frac{\partial G_i}{\partial x^k}, \quad j = 1, \dots, s, k = s+1, \dots, N.$$

Using Equation (4.7) and (4.8),

$$\tilde{X} \log \det A = \sum_{i,j=1}^s A^{ji} \frac{\partial}{\partial x^j} \tilde{X} G_i + \sum_{k=s+1}^N \sum_{j=1}^s \frac{\partial \tilde{X}^k}{\partial x^j} \frac{\partial f_j}{\partial x^k} - \sum_{i=1}^s \frac{\partial \tilde{X}^k}{\partial x^i}.$$

The lemma follows from the above identity and the fact that

$$\sum_{i=s+1}^N \frac{\partial X^i}{\partial x^i} = \sum_{i=s+1}^N \frac{\partial \tilde{X}^i}{\partial x^i} + \sum_{i=s+1}^N \sum_{j=1}^s \frac{\partial \tilde{X}^i}{\partial x^j} \cdot \frac{\partial f_j}{\partial x^i}.$$

□

Go back to the proof of the theorem. Let

$$A = \frac{\partial(F_1^\alpha, \dots, F_s^\alpha)}{\partial(z_{\beta_1}^\alpha, \dots, z_{\beta_s}^\alpha)}$$

and temporary denote $\tilde{X}$ to be the vector field X on $U_\alpha = \{Z_\alpha \neq 0\}$.

The representation of $\tilde{X}$ is in Equation (4.3). Obviously

$$X \log \det \frac{\partial(F_1^\alpha, \dots, F_s^\alpha)}{\partial(z_{\beta_1}^\alpha, \dots, z_{\beta_s}^\alpha)} = \tilde{X} \log \det \frac{\partial(F_1^\alpha, \dots, F_s^\alpha)}{\partial(z_{\beta_1}^\alpha, \dots, z_{\beta_s}^\alpha)}.$$

Using the Lemma 4.1, Equation (4.5) becomes

$$(4.9) \quad \begin{aligned} & \operatorname{div} X + X(f) - (N+1 - d_1 - \dots - d_s)\theta \\ &= \sum_{i \neq \alpha} \frac{\partial \tilde{X}^i}{\partial z_i^\alpha} - \sum_{i=1}^s \sum_{j \in \{\beta_1, \dots, \beta_s\}} A^{ji} \frac{\partial}{\partial z_j^\alpha} \tilde{X} F_i^\alpha \\ & \quad + (N+1 - d_1 - \dots - d_s)(a_{\alpha\alpha} + \sum_{j \neq \alpha} a_{\alpha j} z_j^\alpha). \end{aligned}$$

Since $\sum a_{ii} = 0$, a simple calculation gives

$$(4.10) \quad \sum_{i \neq \alpha} \frac{\partial \tilde{X}^i}{\partial z_i^\alpha} = -(N+1)(a_{\alpha\alpha} + \sum_{j \neq \alpha} a_{\alpha j} z_j^\alpha).$$

Recall the definition of XF_i in Equation (4.1). We see that for $i = 1, \dots, s$,

$$\tilde{X}F_i^\alpha = X \frac{F_i}{Z_\alpha^{d_i}} = \frac{XF_i}{Z_\alpha^{d_i}} - d_i \frac{F_i}{Z_\alpha^{d_i}} (a_{\alpha\alpha} + \sum_{j \neq \alpha} a_{\alpha j} z_j^\alpha).$$

Thus on M_{reg}

$$(4.11) \quad \begin{aligned} & \sum_{i=1}^s \sum_{j \in \{\beta_1, \dots, \beta_s\}} A^{ji} \frac{\partial}{\partial z_j^\alpha} \tilde{X}F_i^\alpha \\ &= \sum_{i=1}^s \sum_{j \in \{\beta_1, \dots, \beta_s\}} A^{ji} Z_\alpha \left(\frac{\frac{\partial}{\partial Z_j} XF_i}{Z_\alpha^{d_i}} - d_i \frac{\frac{\partial F_i}{\partial Z_j}}{Z_\alpha^{d_i}} (a_{\alpha\alpha} + \sum_{j \neq \alpha} a_{\alpha j} z_j^\alpha) \right) \\ &= \kappa - \left(\sum_{i=1}^s d_i \right) (a_{\alpha\alpha} + \sum_{j \neq \alpha} z_j^\alpha), \end{aligned}$$

where we used the fact that on M_{reg} , $F_1 = \dots = F_s = 0$ in the second identity. The theorem follows from Equation (4.9), (4.10) and (4.11).

5. THE COMPUTATION OF THE θ INVARIANTS

Let M be the complete intersection defined in §3. Let $M_0 = CP^N$. $M_k = N_1 \cap \dots \cap N_k$, ($k = 1, \dots, s$). Then $M_s = M$.

We assume that $XF_i = \kappa_i F_i$, $i = 1, \dots, s$.

Let $[Z_0, \dots, Z_N]$ be the homogeneous coordinates of CP^N . Define

$$\xi_i = \frac{|F_i|^2}{\sum_{i=0}^N (|Z_i|^2)^{d_i}}, \quad i = 1, \dots, s.$$

Then ξ_i 's are global smooth functions on CP^N .

In this section, we compute the θ -invariant $\int_M \theta \omega^n$, where for simplifying the notations, we assume that $\omega = \omega_{FS}$ is the Fubini-Study metric of the CP^N . The key result is the following:

Lemma 5.1. *For $k = 2, \dots, s$, we have*

$$(5.1) \quad \int_{M_k} (\theta + \omega)^{N-k+1} = d_k \int_{M_{k-1}} (\theta + \omega)^{N-k+2} + \kappa_k d_1 \cdots d_{k-1},$$

and in addition, we have

$$(5.2) \quad \int_{M_1} (\theta + \omega)^N = \kappa_1.$$

Proof: We have the following identities for $k = 1, \dots, s$

$$(5.3) \quad \begin{aligned} & \bar{\partial} \left(\frac{\sqrt{-1}}{2\pi} \partial \log \xi_k \wedge \theta \omega^{N-k} \right) + i(X) \left(\frac{\partial \log \xi_k}{N-k+1} \omega^{N-k+1} \right) \\ &= -\frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log \xi_k \wedge \theta \omega^{N-k} + \frac{1}{N-k+1} X \log \xi_k \wedge \omega^{N-k+1}. \end{aligned}$$

Integration against M_{k-1} gives

$$\int_{M_{k-1}} \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log \xi_k \wedge \theta \omega^{N-k} = \frac{1}{N-k+1} \int_{M_{k-1}} X \log \xi_k \wedge \omega^{N-k+1}.$$

Since for $k = 1, \dots, s$,

$$\begin{aligned} \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log \xi_k &= [N_k] - d_k \omega \\ X \log \xi_k &= \kappa_k + d_k \theta, \end{aligned}$$

where $[N_k]$ is the divisor of the zero locus of F_k , we have

$$\int_{M_k} \theta \omega^{N-k} - d_k \int_{M_{k-1}} \theta \omega^{N-k+1} = \frac{1}{N-k+1} \int_{M_{k-1}} (\kappa_k + d_k \theta) \omega^{N-k+1}.$$

Thus

$$\int_{M_k} \theta \omega^{N-k} = \frac{N-k+2}{N-k+1} d_k \int_{M_{k-1}} \theta \omega^{N-k+1} + \frac{\kappa_k d_1 \cdots d_{k-1}}{N-k+1}.$$

So (5.1) is proved. To prove (5.2), let's first see that by (5.3),

$$\int_{M_1} \theta \omega^{N-1} = \frac{N+1}{N} d_1 \int_{CP^N} \theta \omega^N + \frac{\kappa_1}{N}.$$

Then (5.2) follows from the simple fact that

$$\int_{CP^N} \theta \omega^N = 0.$$

□

Equation (5.1) can be rewritten as

$$\frac{1}{d_1 \cdots d_k} \int_{M_k} (\theta + \omega)^{N-k+1} = \frac{1}{d_1 \cdots d_{k-1}} \int_{M_{k-1}} (\theta + \omega)^{N-k+2} + \frac{\kappa_k}{d_k}$$

for $k = 2, \dots, s$. Thus we have

Theorem 5.1. *With the notations as above, we have*

$$\int_M \theta \omega^{N-s} = \frac{d_1 \cdots d_s}{N-s+1} \sum_{k=1}^s \frac{\kappa_k}{d_k}.$$

□

Now we prove the main theorem of this paper:

Proof of Theorem 1.1: The theorem follows from Theorem 4.1, Theorem 5.1 and the fact that

$$\omega = (N+1-d_1-\cdots-d_s)\omega_{FS} \in c_1(M).$$

□

6. EXAMPLES

In this section, we use our formula to compute some examples given by Ding-Tian [1], Jeffres [5] and Wu [10]. Recall that, the Futaki invariant defined in [1] and [5] differ from us by a factor 3 in the case of surfaces. So in what follows, the Futaki invariant $F(X)$ is actually three times the Futaki invariant in the previous sections.

Corollary 6.1. *With the notations as in Theorem 1.1, if M is the cubic surface in CP^3 , then the Futaki invariants is*

$$F(X) = -8\kappa.$$

Corollary 6.2. *With the notations as in Theorem 1.1, if M is the variety of the intersection of two quadratic polynomials in CP^4 , then*

$$F(X) = -10(\kappa_1 + \kappa_2).$$

The first four examples are due to Ding and Tian [1]:

Example 1. *Let $X_f \subset CP^3$ be the zero locus of a cubic polynomial f . Put $f = z_0 z_1^2 + z_2 z_3(z_2 - z_3)$, where z_0, z_1, z_2, z_3 are homogeneous coordinates of CP^3 . The X_f has a unique quotient singularity at $p_0 = [1, 0, 0, 0]$. This singularity is of the form C^2/Γ , where Γ is the dihedral subgroup in $SU(2)$ of type D_4 . One can check that X_f is a Kähler orbifold with $c_1(X) > 0$. Let X be the holomorphic vector field whose real part generates the one parameter subgroup $\{\text{diag}(1, e^{3t}, e^{2t}, e^{2t})\}_{t \in \mathbb{R}}$ in $SL(4, \mathbb{C})$. Then X restricts to a holomorphic vector field on X_f and has five zeros $[1, 0, 0, 0]$, $[0, 1, 0, 0]$, $[0, 0, 1, 0]$, $[0, 0, 0, 1]$ and $[0, 0, 1, 1]$.*

We are going to use three methods to compute the Futaki invariants. The first method is the original method in [1].

We can rewrite the function f near $[1, 0, 0, 0]$ in the standard form

$$f = z_1^2 - z_3(z_2^2 - 4z_3^2).$$

In [7] we see that there are standard coverings $C^2 \rightarrow C^2/\Gamma$ by

$$\begin{cases} z_1 = uv(u^4 - v^4) \\ z_2 = u^4 + v^4 \\ z_3 = u^2v^2 \end{cases}.$$

If we assume that on the $u-v$ plane, $X = au\frac{\partial}{\partial u} + bv\frac{\partial}{\partial v}$, then we would have $a = b = \frac{1}{2}$. Since the order of the group D_4 is 8, a computation using the formula in [1, pp 324] shows

$$F(X) = \frac{1}{8} \cdot \frac{1^3}{\frac{1}{4}} + \frac{(-2)^3}{1} + 3\frac{(-1)^3}{-2} = -6.$$

Our second method is a trick which can be generalized to give another prove of the main theorem of this paper in the case of hypersurfaces. Suppose that in $X = au\frac{\partial}{\partial u} + bv\frac{\partial}{\partial v}$, we don't know what a and b is. By using the Bott residue formula, we see that

$$\frac{1}{8} \cdot \frac{(a+b)}{ab} + \frac{(-2)}{1} + 3\frac{(-1)}{-2} = 0$$

and

$$\frac{1}{8} \cdot \frac{(a+b)^2}{ab} + \frac{(-2)^2}{1} + 3\frac{(-1)^2}{-2} = 3.$$

Thus we solved $a = b = \frac{1}{2}$. The Futaki invariant is obtained.

The last method is to use Corollary 6.1, which gives $F(x) = -6$.

Example 2. Let $f = z_0z_1^2 + z_1z_2^2 + z_3^3$ and $X = \{diag(1, e^{6t}, e^{3t}, e^{4t})\}$. Then Corollary 6.1 gives $F(X) = -18$.

Example 3. Let $f = z_0(z_1^2 + z_2^2) + z_3^2z_1$ and $X = diag\{(1, e^{2t}, e^{2t}, e^t)\}$. Then Corollary 6.1 gives $F(X) = -2$.

Example 4. Let $f = z_0(z_1^2 + z_2^2) + z_3^3$. and $X = diag\{(1, e^{3t}, e^{3t}, e^{2t})\}$. Then Corollary 6.1 gives $F(X) = 0$.

The following examples are given by Jeffres [5]: let $[z_0, z_1, w, x, z]$ be the general point in CP^4 .

Example 5. Let

$$\begin{cases} f = z_0z_1 + w^2 + x^2 \\ g = z_1l(w, x) + z^2 \end{cases},$$

where $l(w, x)$ is a linear function of w, x . Let $X = diag\{1, e^{2t}, e^t, e^t, e^{3/2t}\}$. Then $\kappa_1 = -1/5$ and $\kappa_2 = 4/5$. Using Corollary 6.2, $F(X) = -6$.

Example 6.

$$\begin{cases} f = z_0z_1 + z^2 \\ g = z_1^2 + wx \end{cases}$$

and $X = \text{diag}\{(1, e^{2t}, e^{2t}, e^{2t}, e^t)\}$. Then $\kappa_1 = -4/5$ and $\kappa_2 = 6/5$. So the Futaki invariant $F(X) = -4$ using Corollary 6.2.

The following examples are given by Wu [10].

Example 7. Let $M_0 \subset CP^4$ be the zero locus defined by

$$f = z_0 z_1^2 + z_1 z_2^2 + z_3^3 + z_4^3 = 0$$

and $X = \text{diag}(1, e^{6t}, e^{3t}, e^{4t}, e^{4t})$. Then $\kappa = 9/5$ and by Corollary 1.1, $F(X) = -36$.

Example 8. Let $M_0 \subset CP^3$ be the zero locus defined by

$$f = z_3(z_1^2 - z_0 z_2) + z_2^3 = 0$$

and $X = \text{diag}(1, e^t, e^{2t}, e^{4t})$. Then $\kappa = 3/4$ and by Corollary 1.1, $F(X) = -2$.

REFERENCES

- [1] DING, W., AND TIAN, G. Kähler-Einstein metrics and the generalized Futaki invariant. *Invent Math* 110 (1992), 315–335.
- [2] FUTAKI, A. An obstruction to the existence problem of positive Kähler Einstein metrics. *Invent. Math* 73 (1983), 437–443.
- [3] FUTAKI, A. *Kähler-Einstein Metrics and Integral Invariants*, vol. 1314 of *Lect. Notes. Math.* Springer, 1988.
- [4] GRIFFITHS, P., AND HARRIS, J. *Principle of Algebraic Geometry*. John Wiley & Sons, 1978.
- [5] JEFFRES, T. Singular set of some Kähler orbifolds. *Trans. Amer. Math. Soc.* 349(5) (1997), 1961–1971.
- [6] MABUCHI, T. Einstein-Kähler forms, Futaki invariants and convex geometry on toric Fano varieties. *Osaka J. Math* 24 (1987), 705–737.
- [7] PINKHAM, H. Singularities de Klein-I. In *Lect. Notes in Math.* (1976), vol. 777, Springer Verlag, pp. 1–9.
- [8] TIAN, G. The K-energy on hypersurfaces and stability. *Comm. Anal. Geom* 2(2) (1994), 239–265.
- [9] TIAN, G. Kähler-Einstein metrics with positive scalar curvature. *Invent. Math* 137 (1997), 1–37.
- [10] WU, Z. *Kähler-Einstein Metrics and K-Stability*. PhD thesis, Columbia University, New York, NY 10027, May 1998.

(Zhiqin Lu) DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, NY, NY 10027

E-mail address: lu@cpw.math.columbia.edu